Veronica longifolia

Habitus and growth type

Plant height [m]: **0.94**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **19.17**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.07**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **5.1**
Moisture indicator value: **7.4**
Reaction indicator value: **6.6**
Nutrient indicator value: **6.2**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.86**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.68**
Disturbance severity: **0.57**
Disturbance severity (herb layer): **0.49**
Mowing frequency: **0.57**
Grazing pressure: **0.19**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub**

Distribution

Continentality: **7**
Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz