Andryala arenaria

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.6**
Temperature indicator value: **7.9**
Moisture indicator value: **2.6**
Reaction indicator value: **4.3**
Nutrient indicator value: **2.9**
Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.55**
Disturbance frequency (herb layer): **1.46**
Disturbance severity: **0.54**
Disturbance severity (herb layer): **0.3**
Mowing frequency: **0.18**
Grazing pressure: **0.27**
Soil disturbance: **0.18**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FI (TUB) *Helianthemetea guttati***, **QC (CHE) *Chenopodietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz