Arctium lappa subsp. *platylepis*

Habitus and growth type

Plant height [m]: **1.43**
Life span: **Biennial or short-lived**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **17.43**

Fruit, seed and dispersal

Seed mass [mg]: **12.16**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Hypertrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Moisture indicator value: **5.6**
Reaction indicator value: **6.9**
Nutrient indicator value: **8.9**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.61**
Disturbance frequency (herb layer): **1.49**
Disturbance severity: **0.71**
Disturbance severity (herb layer): **0.56**
Mowing frequency: **0.41**
Grazing pressure: **0.17**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QF (ART) *Artemisietea vulgaris***

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz