Campanula sibirica subsp. *parviflora*

Habitus and growth type

Plant height [m]: **0.28**
Life span: **Biennial or short-lived**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **14.98**

Fruit, seed and dispersal

Seed mass [mg]: **0.08**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**
Temperature indicator value: **6.8**
Moisture indicator value: **3.2**
Reaction indicator value: **7.7**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.19**
Disturbance frequency (herb layer): **1.75**
Disturbance severity: **0.46**
Disturbance severity (herb layer): **0.42**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.41**
Grazing pressure: **0.33**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CB (PUB) *Quercetea pubescentis*, DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops, R16 Perennial rocky grassland of Central and South-Eastern Europe, R1B Continental dry grassland (true steppe), R1K Balkan and Anatolian oromediterranean dry grassland**

Constant species of EUNIS habitats: **R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops, R16 Perennial rocky grassland of Central and South-Eastern Europe, R1B Continental dry grassland (true steppe), R1K Balkan and Anatolian oromediterranean dry grassland, S36 Low steppic scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **7**
Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz