# *Dianthus scaber* subsp. *scaber*

### Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

### Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **native**

### Ecology

#### Ellenberg-type indicator values

Temperature indicator value: **5**

#### Disturbance indicator values

Disturbance frequency: **1.1**
Disturbance frequency (herb layer): **1.79**
Disturbance severity: **0.5**
Disturbance severity (herb layer): **0.37**
Mowing frequency: **0.27**
Grazing pressure: **0.39**
Soil disturbance: **0.18**

### Habitat and sociology

#### Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

#### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*