

# *Dianthus viscidus* subsp. *viscidus*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Disturbance indicator values

Disturbance frequency: **0.77**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.41**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.03**

Grazing pressure: **0.33**

Soil disturbance: **0.11**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FO \(ANA\) \*Trifolio anatolici-Polygonetea arenastri\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Constant species of EUNIS habitats: [R1K Balkan and Anatolian oromediterranean dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**