

# *Eragrostis pilosa* subsp. *pilosa*

## Habitus and growth type

Plant height [m]: **0.2**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **42.06**

## Fruit, seed and dispersal

Seed mass [mg]: **0.07**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **7.1**

Reaction indicator value: **6.8**

Nutrient indicator value: **5.8**

Salinity indicator value: **0.2**

### Disturbance indicator values

Disturbance frequency: **1.75**

Disturbance frequency (herb layer): **2.02**

Disturbance severity: **0.81**

Disturbance severity (herb layer): **0.8**

Mowing frequency: **0.37**

Grazing pressure: **0.16**

Soil disturbance: **0.49**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [QD \(DIG\) \*Digitario sanguinalis-Eragrostietea minoris\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V34 Trampled xeric grassland with annuals](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Sparsely vegetated (incl. rock and scree), Synanthropic**

## **Distribution**

Continentality amplitude: **8**