

# *Festuca stricta* subsp. *sulcata*

## Habitus and growth type

Plant height [m]: **0.39**

## Fruit, seed and dispersal

Seed mass [mg]: **0.46**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Ecology

Disturbance indicator values

Disturbance frequency: **1.11**

Disturbance frequency (herb layer): **1.82**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.74**

Grazing pressure: **0.28**

Soil disturbance: **0.17**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) \*Trifolio-Geranietea sanguinei\*](#), [DA \(FES\) \*Festuco-Brometea\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [R1B Continental dry grassland \(true steppe\)](#), [R1C Desert steppe](#), [S36 Low steppic scrub](#)

Constant species of EUNIS habitats: [R16 Perennial rocky grassland of Central and South-Eastern Europe](#), [R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#), [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [R1B Continental dry grassland \(true steppe\)](#), [R1C Desert steppe](#), [S36 Low steppic scrub](#), [T3H Larix light taiga](#)

Dominant species of EUNIS habitats: [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

## Distribution

Continentality: **8**

Continentality amplitude: **4**