

# *Sagina subulata* var. *subulata*

## Habitus and growth type

Plant height [m]: **0.05**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Nutrient indicator value: **2.4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.44**

Disturbance frequency (herb layer): **2.01**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.27**

Grazing pressure: **0.3**

Soil disturbance: **0.26**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Grassland (non-alpine, non-saline)**

### **Distribution**

Continentality: **4**

Continentality amplitude: **4**