Cytinaceae

Habitus and growth type

Plant height [m]: **0.04**
Life span: **Perennial**
Life form: **Geophyte**
Spinescence: **not spinescent**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **endophytic parasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.5**
Temperature indicator value: **9.1**
Moisture indicator value: **2.5**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.42**
Disturbance frequency (herb layer): **1.35**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.27**
Mowing frequency: **0**
Grazing pressure: **0.29**
Soil disturbance: **0.1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S62 Western acidophilous garrigue**

Constant species of EUNIS habitats: **S62 Western acidophilous garrigue**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz