Ceratocarpus

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **4.99**
Dispersal mode: **Anemochory**
Dispersal distance class: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **5**
Moisture indicator value: **3.5**

Disturbance indicator values

Disturbance frequency: **2.15**
Disturbance frequency (herb layer): **2.08**
Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.21**
Grazing pressure: **0.34**
Soil disturbance: **0.29**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***, **EA (LER) *Artemisietea lerchianae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1C Desert steppe**, **S67 Aralo-Caspian semi-**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

desert, **S68 Semi-desert sand dune with sparse scrub**

Constant species of EUNIS habitats: **R1C Desert steppe**, **S67 Aralo-Caspian semi-desert**, **S68 Semi-desert sand dune with sparse scrub**

Dominant species of EUNIS habitats: **S68 Semi-desert sand dune with sparse scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Semi-desert**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz