

Neatostema

Habitus and growth type

Plant height [m]: **0.13**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **16.07**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.23**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **10**

Moisture indicator value: **2**

Reaction indicator value: **7.5**

Nutrient indicator value: **5.5**

Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **1.47**

Disturbance frequency (herb layer): **1.83**

Disturbance severity: **0.73**

Disturbance severity (herb layer): **0.67**

Mowing frequency: **0.2**

Grazing pressure: **0.4**

Soil disturbance: **0.32**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) *Stipo-Trachynietea distachyae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#)

Constant species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub**