Crepis foetida aggr.

Habitus and growth type

Plant height [m]: **0.23**
Life span: **Annual, Biennial or short-lived**
Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **16.08**

Flower

Flowering period: **May-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.61**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.7**
Moisture indicator value: **3.6**
Reaction indicator value: **6.6**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.27**
Disturbance frequency (herb layer): **1.84**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.62**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.55**
Grazing pressure: **0.27**
Soil disturbance: **0.36**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S72 Eastern Mediterranean spiny heath (phrygana)**
Constant species of EUNIS habitats: **S72 Eastern Mediterranean spiny heath (phrygana)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz