Drimia maritima aggr.

Habitus and growth type

Plant height [m]: **1.05**
Life span: **Perennial**
Life form: **Geophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **19.83**

Flower

Flowering period: **August-October**

Fruit, seed and dispersal

Seed mass [mg]: **7.59**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **9.7**
Moisture indicator value: **3**
Reaction indicator value: **7**
Nutrient indicator value: **2.5**
Salinity indicator value: **0.7**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.36**
Disturbance frequency (herb layer): **1.54**
Disturbance severity: **0.71**
Disturbance severity (herb layer): **0.33**
Mowing frequency: **0.08**
Grazing pressure: **0.34**
Soil disturbance: **0.18**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz