# *Chenopodium album* aggr.

### Habitus and growth type

Plant height [m]: **0.95**
Life span: **Annual**
Life form: **Therophyte**

### Leaf

Specific leaf area [$mm^2$/mg]: **17.8**

### Flower

Flowering period: **May-December**

### Fruit, seed and dispersal

Seed mass [mg]: **0.62**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **7.9**
Moisture indicator value: **4.2**
Reaction indicator value: **6.3**
Nutrient indicator value: **6.8**
Salinity indicator value: **0.3**

#### Disturbance indicator values

Disturbance frequency: **1.56**
Disturbance frequency (herb layer): **1.94**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance severity: **0.82**
Disturbance severity (herb layer): **0.81**
Mowing frequency: **0.55**
Grazing pressure: **0.14**
Soil disturbance: **0.71**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: **S68 Semi-desert sand dune with sparse scrub**, **V11 Intensive unmixed crops**, **V12 Mixed crops of market gardens and horticulture**, **V13 Arable land with unmixed crops grown by low-intensity agricultural methods**, **V15 Bare tilled, fallow or recently abandoned arable land**, **V34 Trampled xeric grassland with annuals**, **V37 Annual anthropogenic herbaceous vegetation**
Constant species of EUNIS habitats: **MA221 Atlantic saltmarsh driftline**, **N1E Black Sea broad-leaved coastal dune forest**, **Q61 Periodically exposed shore with stable, eutrophic sediments with pioneer or ephemeral vegetation**, **S68 Semi-desert sand dune with sparse scrub**, **U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region**, **V11 Intensive unmixed crops**, **V12 Mixed crops of market gardens and horticulture**, **V13 Arable land with unmixed crops grown by low-intensity agricultural methods**, **V15 Bare tilled, fallow or recently abandoned arable land**, **V34 Trampled xeric grassland with annuals**, **V35 Trampled mesophilous grassland with annuals**, **V37 Annual anthropogenic herbaceous vegetation**, **V38 Dry perennial anthropogenic herbaceous vegetation**
Dominant species of EUNIS habitats: **V37 Annual anthropogenic herbaceous vegetation**

### Broad habitat

Occurrence in broad habitats: **Wetland, Semi-desert, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

## Distribution

Continentality amplitude: **9**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*