# *Limonium binervosum* aggr.

### Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

### Leaf

Specific leaf area [$mm^2$/mg]: **12.1**

### Fruit, seed and dispersal

Seed mass [mg]: **0.73**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Saline**

#### Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **6**
Reaction indicator value: **7.5**
Nutrient indicator value: **4.5**
Salinity indicator value: **5.8**

#### Disturbance indicator values

Disturbance frequency: **1.54**
Disturbance frequency (herb layer): **1.57**
Disturbance severity: **0.48**
Disturbance severity (herb layer): **0.47**
Mowing frequency: **0.06**
Grazing pressure: **0.27**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Soil disturbance: **0.28**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: **MA251 Mediterranean upper saltmarsh**, **N31 Atlantic and Baltic rocky sea cliff and shore**

Constant species of EUNIS habitats: **MA251 Mediterranean upper saltmarsh**, **N31 Atlantic and Baltic rocky sea cliff and shore**

### Broad habitat

Occurrence in broad habitats: **Coastal cliff**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*