

Linum strictum aggr.

Habitus and growth type

Plant height [m]: **0.33**

Life span: **Annual**

Life form: **Therophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **35.96**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.72**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.1**

Moisture indicator value: **3**

Reaction indicator value: **6.7**

Nutrient indicator value: **2.5**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.91**

Disturbance frequency (herb layer): **1.68**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.55**

Mowing frequency: **0.25**

Grazing pressure: **0.35**

Soil disturbance: **0.24**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1E Mediterranean tall perennial dry grassland](#), [R1F Mediterranean annual-rich dry grassland](#), [S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

Constant species of EUNIS habitats: [R1E Mediterranean tall perennial dry grassland](#), [R1F Mediterranean annual-rich dry grassland](#), [S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**