Taraxacum sect. Scariosa

Habitus and growth type

Plant height [m]: **0.13**
Life form: **Hemicryptophyte**

Flower

Flowering period: **January-April, August-December**

Fruit, seed and dispersal

Seed mass [mg]: **0.73**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry, Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline, Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7.8**
Moisture indicator value: **4.3**
Reaction indicator value: **6.5**
Nutrient indicator value: **4.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.68**
Disturbance frequency (herb layer): **1.81**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.53**
Mowing frequency: **0.26**
Grazing pressure: **0.31**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.32**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S75 Eastern Mediterranean mountain hedgehog-heath**

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz