Aegilops neglecta

Habitus and growth type

Plant height [m]: **0.21**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **23.1**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **20.23**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Moisture indicator value: **2.9**
Reaction indicator value: **7.3**
Nutrient indicator value: **5.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.03**
Disturbance frequency (herb layer): **1.9**
Disturbance severity: **0.68**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.61**
Mowing frequency: **0.52**
Grazing pressure: **0.29**
Soil disturbance: **0.34**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz