# *Agropyron dasyanthum*

### Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

### Fruit, seed and dispersal

Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Mesic**

#### Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **5.5**
Moisture indicator value: **5**

#### Disturbance indicator values

Disturbance frequency: **1.63**
Disturbance frequency (herb layer): **1.66**
Disturbance severity: **0.36**
Disturbance severity (herb layer): **0.36**
Mowing frequency: **0.28**
Grazing pressure: **0.13**
Soil disturbance: **0.42**

### Habitat and sociology

#### Syntaxon

Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea canescentis***

#### EUNIS habitat

Diagnostic species of EUNIS habitats: **R11 Pannonian and Pontic sandy steppe**

#### Broad habitat

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*