Alchemilla straminea

Habitus and growth type

Plant height [m]: **0.19**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **3.5**
Moisture indicator value: **8.3**
Reaction indicator value: **4.3**
Nutrient indicator value: **3.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.58**
Disturbance frequency (herb layer): **1.62**
Disturbance severity: **0.43**
Disturbance severity (herb layer): **0.42**
Mowing frequency: **0.57**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.21**
Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OA (MON) *Montio-Cardaminetea***

Broad habitat

Occurrence in broad habitats: **Spring, Mire, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz