Allium ursinum

Habitus and growth type

Plant height [m]: **0.3**
Life span: **Perennial**
Life form: **Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **37.25**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **6.04**
Dispersal mode: **Myrmecochory, Anthropochory**
Dispersal distance class: **3, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Reaction indicator value: **7.4**
Nutrient indicator value: **7.6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.06**
Disturbance frequency (herb layer): **1.33**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.16**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.01**
Grazing pressure: **0.18**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae*, HA (POP) *Alno glutinosae-Populetea albae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T1F Ravine forest**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz