Allosorus tinaei

Habitus and growth type

Plant height [m]: **0.14**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **9**
Moisture indicator value: **2**
Reaction indicator value: **3**
Nutrient indicator value: **1**
Salinity indicator value: **0**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **KC (ASP) *Asplenietea trichomanis***

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz