Armeria canescens

Habitus and growth type

Plant height [m]: **0.34**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-September**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Moisture indicator value: **3**
Reaction indicator value: **7**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.42**
Disturbance frequency (herb layer): **1.84**
Disturbance severity: **0.39**
Disturbance severity (herb layer): **0.37**
Mowing frequency: **0.19**
Grazing pressure: **0.36**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.15**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R14 Perennial rocky grassland of the Italian Peninsula**, **R1K Balkan and Anatolian oromediterranean dry grassland**, **R45 Alpine and subalpine calcareous grassland of the Balkans and Apennines**

Constant species of EUNIS habitats: **R14 Perennial rocky grassland of the Italian Peninsula**, **R1K Balkan and Anatolian oromediterranean dry grassland**, **R45 Alpine and subalpine calcareous grassland of the Balkans and Apennines**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz