

Astragalus cicer

Habitus and growth type

Plant height [m]: **0.49**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **25.99**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **3.31**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**

Temperature indicator value: **5.9**

Moisture indicator value: **4.3**

Reaction indicator value: **7.6**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.07**

Disturbance frequency (herb layer): **1.71**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.56**

Mowing frequency: **0.78**

Grazing pressure: **0.2**

Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CL \(GER\) *Trifolio-Geranietea sanguinei***](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**

Continentality amplitude: **4**