

Avena sterilis

Habitus and growth type

Plant height [m]: **0.69**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **24.07**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **48.26**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Moisture indicator value: **3.3**

Reaction indicator value: **7.1**

Nutrient indicator value: **5.5**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.19**

Disturbance frequency (herb layer): **1.9**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.77**

Mowing frequency: **0.52**

Grazing pressure: **0.21**

Soil disturbance: **0.5**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [V32 Mediterranean subnitrophilous annual grassland](#), [V33 Dry Mediterranean land with unpalatable non-vernal herbaceous vegetation](#)

Constant species of EUNIS habitats: [V32 Mediterranean subnitrophilous annual grassland](#), [V33 Dry Mediterranean land with unpalatable non-vernal herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**