Ballota acetabulosa

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Seed mass [mg]: **1.7**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **11**
Moisture indicator value: **4**
Reaction indicator value: **7**
Nutrient indicator value: **5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.21**
Disturbance frequency (herb layer): **1.57**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.26**
Mowing frequency: **0.01**
Grazing pressure: **0.32**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S72 Eastern Mediterranean spiny heath (phrygana)**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz