Biscutella didyma

Habitus and growth type

Plant height [m]: **0.45**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Flower

Flowering period: **January-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.77**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**
Temperature indicator value: **9.8**
Moisture indicator value: **3**
Reaction indicator value: **6.8**
Nutrient indicator value: **2.7**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.6**
Disturbance frequency (herb layer): **1.76**
Disturbance severity: **0.76**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.48**
Mowing frequency: **0.17**
Grazing pressure: **0.33**
Soil disturbance: **0.29**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S63 Eastern garrigue**, **S72 Eastern Mediterranean spiny heath (phrygana)**
Constant species of EUNIS habitats: **S63 Eastern garrigue**, **S72 Eastern Mediterranean spiny heath (phrygana)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz