# *Brassica repanda*

## Habitus and growth type

Plant height [m]: **0.16**
Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

## Leaf

Specific leaf area [$mm^2$/mg]: **9.89**

## Flower

Flowering period: **March-August**

## Fruit, seed and dispersal

Seed mass [mg]: **1.4**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.5**
Moisture indicator value: **3**
Reaction indicator value: **7**
Nutrient indicator value: **3**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.17**
Disturbance frequency (herb layer): **1.62**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance severity: **0.44**
Disturbance severity (herb layer): **0.39**
Mowing frequency: **0.12**
Grazing pressure: **0.28**
Soil disturbance: **0.2**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: **S73 Western Mediterranean mountain hedgehog-heath**

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*