

Bromopsis cappadocica

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **5.5**

Moisture indicator value: **4**

Disturbance indicator values

Disturbance frequency: **0.76**

Disturbance frequency (herb layer): **1.66**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.25**

Grazing pressure: **0.3**

Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S75 Eastern Mediterranean mountain hedgehog-heath](#)

Constant species of EUNIS habitats: [S75 Eastern Mediterranean mountain hedgehog-heath](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),**

Oromediterranean scrub, Forest, Sparsely vegetated (incl. rock and scree)