

# *Bromopsis erecta*

## Habitus and growth type

Plant height [m]: **0.55**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **19.24**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **4.9**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **5.3**

Moisture indicator value: **3.1**

Reaction indicator value: **7.9**

Nutrient indicator value: **2.9**

Salinity indicator value: **0.2**

### Disturbance indicator values

Disturbance frequency: **0.88**

Disturbance frequency (herb layer): **1.75**  
Disturbance severity: **0.52**  
Disturbance severity (herb layer): **0.43**  
Mowing frequency: **0.58**  
Grazing pressure: **0.28**  
Soil disturbance: **0.17**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#), [R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe](#), [R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#), [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Constant species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#), [R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe](#), [R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#), [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [R1H Iberian oromediterranean basiphilous dry grassland](#), [R32 Mediterranean short moist grassland of lowlands](#), [R45 Alpine and subalpine calcareous grassland of the Balkans and Apennines](#), [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#), [S31 Lowland to montane temperate and submediterranean Juniperus scrub](#), [S52 Submediterranean pseudomaquis](#), [S53 Spartium junceum scrub](#), [S73 Western Mediterranean mountain hedgehog-heath](#), [T1A Mediterranean thermophilous deciduous forest](#), [T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest](#)

Dominant species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#), [R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe](#), [R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#), [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [S53 Spartium junceum scrub](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**, **Alpine-subalpine and arctic grassland**, **Oromediterranean scrub**, **Scrub**, **Forest**, **Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**  
Continentality amplitude: **4**