Bupleurum semicompositum

Habitus and growth type

Plant height [m]: **0.18**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.2**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **11**
Moisture indicator value: **4.5**
Reaction indicator value: **9**
Nutrient indicator value: **3**
Salinity indicator value: **6**

Disturbance indicator values

Disturbance frequency: **1.4**
Disturbance frequency (herb layer): **1.51**
Disturbance severity: **0.65**
Disturbance severity (herb layer): **0.63**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.13**
Grazing pressure: **0.32**
Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FJ (TRA) *Stipo-Trachynietea distachyae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R61 Mediterranean inland salt steppe**, **S66 Mediterranean halo-nitrophilous scrub**

Constant species of EUNIS habitats: **S66 Mediterranean halo-nitrophilous scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz