

Callitriche stagnalis

Habitus and growth type

Plant height [m]: **0.34**

Life span: **Annual**

Life form: **Hydrophyte, Therophyte**

Leaf

Specific leaf area [mm²/mg]: **32.23**

Flower

Flowering period: **March-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.12**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **6.5**

Moisture indicator value: **10.6**

Reaction indicator value: **5.3**

Nutrient indicator value: **4.2**

Salinity indicator value: **0.6**

Disturbance indicator values

Disturbance frequency: **1.33**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.32**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.07**

Grazing pressure: **0.07**

Soil disturbance: **0.08**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [NB \(POT\) Potamogetonetea](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

Distribution

Continentality: **4**

Continentality amplitude: **4**