Callitriche stagnalis

Habitus and growth type

Plant height [m]: **0.34**
Life span: **Annual**
Life form: **Hydrophyte, Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **32.23**

Flower

Flowering period: **March-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.12**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **6.5**
Moisture indicator value: **10.6**
Reaction indicator value: **5.3**
Nutrient indicator value: **4.2**
Salinity indicator value: **0.6**

Disturbance indicator values

Disturbance frequency: **1.33**
Disturbance frequency (herb layer): **1.67**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.32**
Disturbance severity (herb layer): **0.3**
Mowing frequency: **0.07**
Grazing pressure: **0.07**
Soil disturbance: **0.08**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **NB (POT) *Potamogetonetea***

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz