

Campanula latifolia

Habitus and growth type

Plant height [m]: **0.83**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **55.58**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.27**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.8**

Temperature indicator value: **4.8**

Moisture indicator value: **6.1**

Reaction indicator value: **7.7**

Nutrient indicator value: **6.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.12**

Disturbance frequency (herb layer): **1.35**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.2**

Mowing frequency: **0.03**

Grazing pressure: **0.18**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) *Carpino-Fagetea sylvaticae*](#),
[CS \(MUL\) *Mulgedio-Aconitetea*](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Alpine-subalpine and arctic grassland, Scrub, Forest, Synanthropic**

Distribution

Continentality: **5**

Continentality amplitude: **4**