Campanula radicosa

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.4**
Temperature indicator value: **5.5**
Moisture indicator value: **5.8**
Reaction indicator value: **6.3**
Nutrient indicator value: **3.6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.31**
Disturbance frequency (herb layer): **2.31**
Disturbance severity: **0.33**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0.02**
Grazing pressure: **0.34**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FO (ANA) *Trifolio anatolici-Polygonetea arenastri***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1K Balkan and Anatolian oromediterranean dry grassland**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Constant species of EUNIS habitats: **R1K Balkan and Anatolian oromediterranean dry grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz