Carex fimbriata

Habitus and growth type

Plant height [m]: **0.25**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **2.3**
Moisture indicator value: **4.3**
Reaction indicator value: **2**
Nutrient indicator value: **1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.16**
Disturbance frequency (herb layer): **1.39**
Disturbance severity: **0.29**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.06**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.28**
Soil disturbance: **0.09**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CT (TRI) *Juncetea trifidi***

Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz