

Cassiope tetragona

Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Chamaephyte, Dwarf shrub**

Leaf

Specific leaf area [mm²/mg]: **5.68**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Moisture indicator value: **5**

Disturbance indicator values

Disturbance frequency: **1.65**

Disturbance frequency (herb layer): **1.65**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0**

Grazing pressure: **0.3**

Soil disturbance: **0.09**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**AB \(KOB\) Carici rupestris-Kobresietea bellardii**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R42 Boreal and Arctic acidophilous alpine grassland](#), [S11 Shrub tundra](#), [U52 Polar desert](#)

Constant species of EUNIS habitats: [R42 Boreal and Arctic acidophilous alpine grassland](#), [S11 Shrub tundra](#), [U52 Polar desert](#)

Dominant species of EUNIS habitats: [S11 Shrub tundra](#)

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**