Catabrosa aquatica

Habitus and growth type

Plant height [m]: **0.29**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **44.05**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.37**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **4.3**
Moisture indicator value: **9.3**
Reaction indicator value: **6.7**
Salinity indicator value: **0.9**

Disturbance indicator values

Disturbance frequency: **1.67**
Disturbance frequency (herb layer): **1.77**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.31**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0.14**
Grazing pressure: **0.16**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OD (PHR) *Phragmito-Magnocaricetea***

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Grassland (non-alpine, non-saline)**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz