Centaurea cuneifolia

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.4**
Temperature indicator value: **8.4**
Moisture indicator value: **3.5**

Disturbance indicator values

Disturbance frequency: **1.28**
Disturbance frequency (herb layer): **1.59**
Disturbance severity: **0.35**
Disturbance severity (herb layer): **0.33**
Mowing frequency: **0.11**
Grazing pressure: **0.28**
Soil disturbance: **0.25**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**
Constant species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz