Cerastium pumilum

Habitus and growth type

Plant height [m]: **0.11**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **22.56**

Flower

Flowering period: **February-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.08**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Moisture indicator value: **2.3**
Reaction indicator value: **7.3**
Nutrient indicator value: **2.1**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.56**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.99**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.5**
Mowing frequency: **0.39**
Grazing pressure: **0.35**
Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CK (SED) *Sedo-Scleranthetea*, FJ (TRA) *Stipo-Trachynietea distachyae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R12 Cryptogam- and annual-dominated vegetation on siliceous rock outcrops, R13 Cryptogam- and annual-dominated vegetation on calcareous and ultramafic rock outcrops, S34 Balkan-Anatolian submontane genistoid scrub**

Constant species of EUNIS habitats: **R12 Cryptogam- and annual-dominated vegetation on siliceous rock outcrops, R13 Cryptogam- and annual-dominated vegetation on calcareous and ultramafic rock outcrops, R1F Mediterranean annual-rich dry grassland, S34 Balkan-Anatolian submontane genistoid scrub**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz