Cerastium soleirolii

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.27**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **4**
Moisture indicator value: **5**
Reaction indicator value: **3**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.05**
Disturbance frequency (herb layer): **1.54**
Disturbance severity: **0.35**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.02**
Grazing pressure: **0.24**
Soil disturbance: **0.09**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FQ (GEN) *Carici-Genistetea lobelii***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1J Cyrno-Sardean oromediterranean siliceous dry grassland**, **S74 Central Mediterranean mountain hedgehog-heath**, **U24 Mediterranean siliceous scree**

Constant species of EUNIS habitats: **R1J Cyrno-Sardean oromediterranean siliceous dry grassland**, **S74 Central Mediterranean mountain hedgehog-heath**, **U24 Mediterranean siliceous scree**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz