Cerinthe minor

Habitus and growth type

Plant height [m]: **0.37**
Life span: **Annual, Biennial or short-lived, Perennial**
Life form: **Hemicryptophyte, Therophyte**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **9.06**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**
Moisture indicator value: **4.3**
Reaction indicator value: **6.8**
Nutrient indicator value: **4.2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.01**
Disturbance frequency (herb layer): **1.55**
Disturbance severity: **0.59**
Disturbance severity (herb layer): **0.52**
Mowing frequency: **0.55**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.22**
Soil disturbance: **0.29**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **7**
Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz