Clinopodium graveolens

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.39**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Temperature indicator value: **11**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.86**
Disturbance frequency (herb layer): **1.67**
Disturbance severity: **0.44**
Disturbance severity (herb layer): **0.37**
Mowing frequency: **0.21**
Grazing pressure: **0.32**
Soil disturbance: **0.19**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**, **T38 Mediterranean montane Cedrus forest**, **T3D Mediterranean Cupressaceae forest**
Constant species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**, **T38 Mediterranean montane Cedrus forest**, **T3D Mediterranean**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Cupressaceae forest

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz