Cochlearia glastifolia

Habitus and growth type

Plant height [m]: **1.05**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **7**
Moisture indicator value: **4**
Reaction indicator value: **6**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz