Cochlearia groenlandica

Habitus and growth type

Life span: **Biennial or short-lived, Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Ellenberg-type indicator values

Moisture indicator value: **8**

Disturbance indicator values

Disturbance frequency: **1.77**
Disturbance frequency (herb layer): **1.77**
Disturbance severity: **0.44**
Disturbance severity (herb layer): **0.44**
Mowing frequency: **0.12**
Grazing pressure: **0.22**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **AE (COC) *Saxifrago cernuae-Cochlearietea groenlandicae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **MA211 Arctic coastal saltmarsh**

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz