Coleanthus subtilis

Habitus and growth type

Plant height [m]: **0.04**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **May-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.03**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Reaction indicator value: **3.8**
Nutrient indicator value: **3.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.99**
Disturbance frequency (herb layer): **1.99**
Disturbance severity: **0.8**
Disturbance severity (herb layer): **0.8**
Mowing frequency: **0**
Grazing pressure: **0.19**
Soil disturbance: **0.39**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OC (ISO) *Isoëto-Nanojuncetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q62 Periodically exposed shore with stable, mesotrophic sediments with pioneer or ephemeral vegetation**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

Distribution

Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz