Conioselinum tataricum

Habitus and growth type

Plant height [m]: **0.8**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.8**
Temperature indicator value: **3.9**
Moisture indicator value: **6.4**
Reaction indicator value: **8**
Nutrient indicator value: **7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.26**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.64**
Disturbance severity (herb layer): **0.3**
Mowing frequency: **0.1**
Grazing pressure: **0.18**
Soil disturbance: **0.13**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BC (BRA) *Brachypodio pinnati-Betuletea pendulae***, **CM (MOL) *Molinio-Arrhenatheretea***, **CS (MUL) *Mulgedio-Aconitetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T3H Larix light taiga**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz