Corydalis solida

Habitus and growth type

Plant height [m]: **0.16**
Life span: **Perennial**
Life form: **Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **43.65**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.95**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **3.7**
Temperature indicator value: **5.9**
Moisture indicator value: **5.5**
Reaction indicator value: **7**
Nutrient indicator value: **6.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.06**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.33**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.16**
Mowing frequency: **0.01**
Grazing pressure: **0.18**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T33 Mediterranean mountain Abies forest**, **T38 Mediterranean montane Cedrus forest**
Constant species of EUNIS habitats: **T38 Mediterranean montane Cedrus forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz