Crepis zacintha

Habitus and growth type

Plant height [m]: **0.21**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **27.55**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **9.7**
Moisture indicator value: **4.5**
Reaction indicator value: **7**
Nutrient indicator value: **5**
Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **1.16**
Disturbance frequency (herb layer): **1.86**
Disturbance severity: **0.75**
Disturbance severity (herb layer): **0.66**
Mowing frequency: **0.33**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.32**
Soil disturbance: **0.36**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz